Abstract

Of the several processes that purportedly contribute to psychological flexibility, that of enhancing self-as-context, or

Comparing a Brief Self-as-Context Exercise to Control-Based and Attention-Placebo Protocols for Coping with Induced Pain

Over the last decade an increasingly large body of clinical and research literature has been accumulated concerning acceptance and commitment therapy (ACT; Hayes, Strosahl, & Wilson, 2012). ACT is a transdiagnostic approach based on a unified model of human functioning W K D W S U R P R W H V S V \ F K R O R J L F D O I O H [L E L O L W \ D V [3] W

As recently detailed by Levin and his colleagues (2012), laboratory-based studies have documented significant positive effect sizes for components within ACT that ostensibly target each of the processes within the hexaflex with the exception of self-as-context. This absence results not from any negative findings that have emerged from investigating ACT-related procedures and exercises designed to enhance self-as-context, but from a general absence of any laboratory-based research that has attempted to isolate components that purportedly foster this process. Self-as-context within ACT is defined as witnessing private events from a particular vantage point (Hayes, Strosahl, & Wilson, 1999) and as being integral to perspective taking (McHugh & Stewart, 2012). In particular, it is the perspective from which we are capable of observing the continual flow of thoughts, sensations, feelings, and other private events, while maintaining a distinction between sensing and what is sensed (Hayes, Strosahl, & Wilson, 1999). The behavior of "HHLQJ WKDW, DP V HHLQJ contributes to psychological flexibility as it promotes contact with the present moment and provides a context for defusion from thoughts and other private events that may otherwise occasion experiential control (Hayes, 1984; Hayes et al., 2012).

It seems pertinent to note that while self-as-context has most often been conceptualized in a somewhat limited way as transcendent perspective taking (Hayes, Strosahl, & Wilson, 1999), it has recently been viewed more expansively as situated within a wider array of related psychological experiences; such as empathy, compassion, and self-compassion; that are thought to emerge from similar types of verbal-social interactions (i.e., deictic relational framing; Hayes et al., 2012).

Method

Participants. Participants included college students enrolled in psychology courses who received extra credit for participation. We first prescreened them through an online system with the Acceptance and Action Questionnaire (AAQ; Hayes et al., 2004). The AAQ is a self-report LQYHQWRU\ RI SV\FKRORJLFDO LQIOH[LELOLW\ FRQVLVWLQJ RI DQ[LRXV , DP XQDEOH WR WDNH FDUH-point Likert scale. Total scores LQ VLELOL range from 9±63, with higher scores reflecting higher levels of psychological inflexibility. Although an updated version of the AAQ (i.e., AAQ-II; Bond et al., 2011) is now available, it had not yet appeared in print when we began this project. In addition, we opted to use the original AAQ because our familiarity with it in screening participants for similar projects

Students who received AAQ scores • 26 were invited to participate in this study, with 122 agreeing to do so. This cut-off score falls within one standard deviation of the AAQ mean and was selected based on previous research indicating that individuals with greater levels psychological flexibility (i.e., lower AAQ scores) tend to display higher pain tolerance (Zettle et al., 2005). Accordingly, we used this criterion to minimize any potential ceiling effects

> The purpose of this strategy is not to eliminate or somehow control your pain, but rather to see if it is possible to change your experience of pain by viewing it from a different perspective as something that is apart from you.

S-AC was based on the observer exercise presented in Hayes, Strosahl, and Wilson (1999, pp. 193-195), but was altered slightly by adding the following segment:

> Notice that unpleasant experiences, such as pain, that you may struggle with and try to change are not you anyway. No matter how this struggle turns out, you will be there, unchanged. See whether you can take advantage of this connection to let go just a little bit of whatever unpleasant sensations and experiences you may struggle with.

Control-based protocol (CB). We introduced this protocol by saying the following:

> The purpose of this strategy is to introduce you to several techniques that can be used in managing and controlling pain. These techniques may represent an extension and refinement of several things you may already have learned how to do in coping with pain.

CB was developed based on a traditional cognitive-behavioral approach to pain provided by Turk et al. (1983) and presented several strategies designed to help participants control and modify thoughts and emotions related to the experience of pain. Specifically, participants were instructed in and encouraged to use relaxation techniques, such as deep breathing, cognitive restructuring techniques, such as self-talk, and positive imagery.

(pre-post) ANOVAs on each of the variables. The only significant effects that were detected were for measurement occasion for threshold, $\lambda = .87, F(1,63) = 9.64, p < .01, \eta_p^2 = .13$; tolerance, $\lambda = .73, F(1,63) = 23.9, p < .01, \eta_p^2 = .28$, and endurance, $\lambda = .77, F(1,63) = 18.06, p < .01, \eta_p^2 = .22$, suggesting a possible placebo effect and/or demand characteristic in which all three protocols were equally efficacious in improving the ability of participants to cope with pain. Of these two possibilities, an equivalence in demand characteristics would seem the less probable, given that the expressed SXUSRVHV RI EROWK &% ³PDQDJLQJ DQG RYHUFRPLQJ SDLQ´ DEOH WR PDQDJH SDLQ´ DQG ³HQGXUDQFH´ DV WKH RXWFRPH YDULDEOH ³WKUHVKROG´ DQG ³WROHUDQFH´ DV WKH RXWFRPH YDULDEOHV However, if any such differential demand characteristics were present, they were not reflected in the value and utility ratings.

Alternatively, WKH RYHUDOO ILQGLQJV FRXOG DOVR UHIOHFW that the protocols were inert and participants simply became more tolerant of the pain upon experiencing it a second time. In order to further explore this possibility, we conducted a series of paired-samples, one-tailed t-tests within each of the protocol conditions. As can be seen in Table 3, all of the protocols significantly increased pain tolerance, while only CB and AP increased pain threshold and endurance, suggesting that the overall findings could not be simply attributed to either a practice or placebo effect. None of the protocols led to significant reductions in pain intensity. This particular finding is not that unusual as other studies have reported similar results for ostensibly active interventions. For instance, Sharpe et al. (2010) found that brief relaxation and metacognitive attention training conditions were equally ineffective in decreasing the intensity of cold pressor-induced pain.

Another lens through which we viewed differences in the relative impact of the three protocols in increasing pain tolerance involved a comparison of effect sizes. Within-condition effect sizes reported in Table 2 were compu

Because the collective findings of Study 1, as well as effect size comparisons to those of Hayes, Bissett, et al., (1999) adequately established the relative impact of the acceptance-based and attention-placebo conditions, we opted to only present participants in this second study with a contextualized, pain-specific version of the observer exercise.

Method and Procedure

We followed the same procedures and selection criteria for recruiting and screening participants as used in Study 1. A total of 60 college students were initially screened into the study and agreed to participate in until we obtained a final sample size (i.e., N = 22) equal to that within each of the three conditions of Study 1. As can be seen in Table 2

It seems less clear to us why the large effect size for C-SAC o

SELF

selecting a self-as-context–related component that could be presented in this manner. However, more could have been done as our assistants were not blind to protocol assignment in administering the postintervention cold pressor, the manipulation and treatment integrity checks, and the value and utility scales. It would be desirable in future research to further minimize potential bias effects by ensuring that those who interact with participants remain blind to the protocols presented to them.

The most obvious limitations to external validity are hardly unique to our project, but apply to analogue research in general. As with the majority of other laboratory component studies (Levin et al., 2012), our participants were college students who may not be representative of other nonclinical populations. Consequently, it may be useful to determine if our overall findings are replicable with community samples that may differ from ours in age and educational background, for example. Perhaps a more germane concern regarding the generalizability of our findings is whether they would extend to presenting a contextualized version of the observer exercise to those faced with other psychological and biological challenges. For instance, because pain is a more ubiquitous experience, contextualizing the observer exercise for use with the cold pressor may prove to be more viable than modifying it to assist participants about to encounter induced panic-like symptoms (e.g., Feldner et al., 2003).

The ability to generalize our overall findings to clinical populations who struggle with pain more generally, and chronic pain, in particular, may also be restricted. Pain induced by the cold pressor is more analogous to acute pain inherent in certain medical procedures, such as injections and changing of wound/burn dressings. Traditional cognitive behavioral therapies that incorporate many of the same strategies included in the CB protocol of Study 1; such as relaxation techniques, imagery, and use of coping self-statements, represent an empirically

supported approach for acute, procedure-related pain (e.g., Powers, 1999). While cognitive-behavioral interventions have also been shown to be efficacious in targeting chronic pain (Morley, Eccleston, & Williams, 1999), those receiving ACT have reported greater satisfaction with their treatment (Wetherell et al., 2011). The degree to which ACT for chronic pain is appreciably enhanced by components focused on self-as-context, such as the observer exercise, however, is an empirical question that cannot be adequately addressed merely by extending our findings, but only through further research.

A final limitation on the external validity of our findings that is more specific to our project and not endemic to analogue research more broadly concerns the relatively higher proportion of participants (i.e., 30% of the total participant pool) who were excluded for failing the manipulation and treatment integrity checks of Study 2. Participants in general found it easier to understand and apply the control-based protocol than those dealing with self-as-context, and had relatively more difficulty with S-SAC than C-SAC. On first consideration, such findings might suggest that control-based therapeutic approaches to pain management may have wider applicability and utility than those, such as ACT, that seek to enhance flexible perspective taking. The strong empirical support that ACT has received in treatment of chronic pain, however, would argue against such an interpretation, as would we believe some further reflection on how the observer exercise was presented within our two studies as compared to how it is introduced and presented clinically.

Because of the cultural dominance of the experiential control agenda and the counterintuitive nature of the perspective the observer exercise seeks to foster, it does not seem surprising that college students, particularly those less fluent in English as suggested by some of the findings of Study 2, would find grasping and applying it to be a challenge. However, when

SELF-AS-

components suggested by the psychological flexibility model: A meta-analysis of laboratory-based component Studies. *Behavior Therapy, 35*, 741-756. doi:10.1016/j.beth.2012.05.003

Lillis, J., & Hayes, S. C. (2008). Measuring avoidance and inflexibility in weight related problems. *International Journal of Behavioral Consultation and Therapy, 4*, 30-40.

Lillis, J., Hayes, S. C., Bunting, K., & Masuda, A. (2009). Teaching acceptance and mindfulness to improve the lives of the obese: A preliminary test of a theoretical model. *Annals of Behavioral Medicine, 37*, 58-69. doi: 10.1007/s12160-009-9083-x

Loeser,, J. D., & Melzack, R. (1999). Pain: An overview. *The Lancet, 353*, 1607-1609.

McHugh, L., & Stewart, I. (2012). *The self and perspective taking: Contributions and applications from modern behavioral science*. Oakland, CA: Context Press.

Morley, S., Eccleston, C., & Williams, A. (1999). A systematic review and meta-analysis of randomized controlled trials of cognitive behaviour therapy and behaviour therapy for chronic pain in adults, excluding headache. *Pain, 80*, 1-13.

Orsillo, S. M., & Batten, S. J. (2005). Acceptance and commitment therapy in the treatment of posttraumatic stress disorder. *Behavior Modification, 29*, 95-129. doi: 10.1177/0145445504270876

Powers, S. W. (1999). Empirically supported treatments in pediatric psychology: Procedure related pain. *Journal of Pediatric Psychology, 24*, 131-145.

Sharpe, L., Perry, K. N., Rees, G., Dear, B. F., Nicholas, M. K., & Refshauge, K. (2010). A comparison of the effect of attention training and relaxation on responses to pain. *Pain, 150*, 469-476. doi:10.1016/j.pain.2010.05.027

Streff, A., Kuehl, L. K., Michaux, G., & Anton, F. (2010). Differential physiological effects

during tonic painful hand immersion tests using hot and ice water. *European Journal of Pain, 14*, 266-272. doi:10.1016/j.ejpain.2009.05.011

Turk D. C., Meichenbaum D., & Genest M. (1983). *Pain and behavioral medicine. A cognitive-behavioral perspective*. New York: Guilford.

Walser, R., & Westrup, D. (2007). *Acceptance and commitment therapy for the treatment of post-traumatic stress disorder and trauma-related problems: A practitioner's guide to using mindfulness and acceptance strategies*. Oakland, CA: New Harbinger.

Westin, V., Hayes, S. C., & Andersson, G. (2008). Is it the sound or your relationship to it? The role of acceptance in predicting tinnitus impact. *Behaviour Research and Therapy, 46*, 1259-1265. doi: 10.1016/j.brat.2008.08.008

Wetherell, J. L., Afari, N., Rutledge, T., Sorrell, J. T., Stoddard, J. A., Petkus, A. J., . . . Atkinson, J. H. (2011). A randomized, controlled trial of acceptance and commitment therapy and cognitive-behavioral therapy for chronic pain. *Pain, 152*

Sudarijanto, R. P. (2005). Differential strategies in coping with pain as a function of level of experiential avoidance. *The Psychological Record*, *55*, 511-524.

Footnotes

[1]Copies of CDs used in both studies may be obtained by contacting the first author.

Table 1

Means and Standard Deviations for Value and Utility Scale Ratings of Protocols

	Value		Utility	
Protocol	*M*	*SD*	*M*	*SD*
1. G-SAC[a]	6.15	4.13	6.41	3.62
2. CB	8.00	2.66	8.13	2.91
3. AP	5.80	3.32	6.02	3.74
4. C-SAC[b]	8.68	3.76	7.66	3.76
Statistical Analyses				
F	3.51		1.77	
p	0.02		0.16	
n_p^2	.11		.06	
Comparisons	“		N/A	

[a]Denotes generic self-as-context protocol of Study 1.
[b]Denotes contextualized self-as-context protocol of Study 2

Table 2

Characteristics of Participants in the Two Studies

Variable	Study 1(N = 66)		Study 2(N = 22)	
	M	*SD*	*M*	*SD*
AAQ	35.36	6.37	33.00	4.32
Age	21.58	5.79	21.27	4.14
	N	%	N	%
Ethnicity				
White	38	58	13	59
Black	10	15	1	4